

ISOLATION - AT - A - GLANCE

A Airborne Precautions C Contact Precautions CP Contact Precautions Plus D Droplet Precautions S Standard Precautions P Private Room NP Negative Pressure Room	Post appropriate Precaution sign at patient room Cohort patients with the same illness under certain circumstances - (contact Infection Prevention x 8232, 8223, 6961) Transport - notify all receiving parties of isolation status
hand hygiene saves lives hand hygiene saves lives hand hygiene saves lives hand hygiene saves lives hand hygiene saves lives hand hygiene saves lives hand hygiene saves lives hand hygiene saves lives	

CONDITION/ DISEASE	PRECAUTIONS		ROOM	EMPLOYEE PPE	PATIENT PPE for TRANSPORT
	TYPE	DURATION			
MDRO (multi-drug resistant organism)					
Resistant Acinetobacter	C	Lifelong	P	Gown, Gloves	Clean pt gown
ESBL producing organisms	C	Lifelong	P	Gown, Gloves	Clean pt gown
MRSA	C	For 2 years after last positive culture	P	Gown, Gloves	Clean pt gown
VRE	C	Lifelong	P	Gown, Gloves	Clean pt gown
Chickenpox (Varicella Zoster)	A, C	Until all lesions are crusted	NP	Surgical mask (susceptible HCWs should not enter room if immune HCWs are available)	Surgical mask, Clean pt gown
Gastroenteritis - <i>C.difficile</i> toxin	CP	Duration of admission	P	Gown, Gloves	Clean pt gown
Influenza	D	Duration of illness	P	Surgical mask	Surgical mask
Lice	C	24 hrs after start of effective therapy	P	Gown, Gloves	Clean pt gown
Meningitis - Bacterial r/o meningococcal disease	D	24 hrs after start of effective therapy	P	Surgical mask	Surgical mask
Meningitis - Bacterial r/o <i>H. influenzae</i> type b		25 hrs after start of effective therapy	P	Surgical mask	Surgical mask
Meningitis - Viral, Aseptic	S			PPE in accordance with Standard Precautions	
Pertussis	D	Until 5 days after start of effective therapy	P	Surgical mask	Surgical mask
Scabies	C	24hrs after start of effective therapy	P	Gown, Gloves	Clean pt gown
Shingles (Herpes zoster) (varicella-zoster)					
Localized in pt with intact immune system lesions can be contained/covered	C	Duration of illness	P	Gown, Gloves (susceptible HCWs should not provide pt care when other immune HCWs are available)	Clean pt gown
Disseminated disease in any patient. Localized disease in immunocompromised pt until dissemination infection ruled out	A, C	Duration of illness	NP	Surgical mask, Gown, Gloves (susceptible HCWs should not enter room if immune HCWs are available)	Surgical mask
Tuberculosis	A	Contact Infection Prevention Dept.	NP	N95 mask	Surgical mask